

Baljit Dheansa FRCS MSc FRCS(Plast) Consultant Plastic, Cosmetic & Reconstructive Surgeon

My Plastic Surgeon Limited
The McIndoe Centre
Holtye Road
East Grinstead
West Sussex
RH10 3FR

Patient:	Procedure:	Breast Augmentation	On Date:	
RISKS / Concernion	OMPLICATIONS (rare or unusual comp	lications can occur and m	nay not be listed)	
Allergic	reactions to any drugs, antiseptics or	Dehiscence - wound o	pening up - may require	
equipment used		prolonged dressings		
Asymmetry (this may be better or worse than pre-		Skin loss/necrosis – may require prolonged		
operatively)		dressings or surgery including skin grafts		
Bleeding – may require further procedure		Swelling, Inflammation - may take months to		
		resolve		
Bruising – may take weeks to resolve		Slow healing		
Haematoma (collection of blood under skin) – may		Poor scarring - Keloid, Hypertrophic, unsightly -		
require further procedure		may be difficult to treat o	may be difficult to treat or improve	
Infection - may require antibiotics or surgery or		Numbness around or beyond the area treated -		
may cause tissue loss		may increase risk of inadvertent injury in future		
Pain – m	nay be long term	Pulmonary Embolus or	Deep Vein Thrombosis -	
		clots in the lung or legs		
Changes	s in contour of tissues - may be visually	Change over time - tiss	sues may become thinner,	
apparen	t	droop or lose the effect of	of surgery	
Cassific to				
Specific to		ALCI may dayalan thi	io concer io uncommon	
	or capsular contracture – breast shape		is cancer is uncommon –	
	ange or distort – may require surgery or of implant	may need surgery and/or	chemotherapy	
	•	Implant may runtura furt	ther ourgen, will be needed	
	nipple sensation – or increased sensitivity		ther surgery will be needed	
_	or rippling of the implant - may require		the implant - may need	
Strotch	morke may ennour	drainage or surgery Fold under breast may be	lower	
	marks may appear			
	will need to be removed if infected		e almost certainly required	
	ement or movement of the implant post-	be detected in breast milk	be affected – silicone may	
	ely – may require surgery	Cancer detection may be		
	ize may increase of implant below or above muscle may		pregnancy, weight gain or	
	risks – weakness, bleeding, visible	loss	pregnancy, weight gain of	
	ent may be affected	1055		
	intment with final appearance	No guarantee of size im	plant size may need to be	
Ызарро	пипен мин ппагарреагансе	modified according to tiss		
Dheansa's complication science. It is may affect I permit the publication	a has discussed the above risks and com Breast Augmentation information leaflet [ons can occur but may be less common as understand the risks, benefits and limitatio my ability to work, socialise or do certain a ne use of my non-identifiable photograpl and to be shown to other patients. I can v	(Please Tick). I also und the practice of medicine at the practice of medicine at the proposed proceduractivities. This may sometimes (pre-and post-operative)	derstand other risks and and surgery is not an exact ure. Certain complications nes be long term.	
Patient Si	gnature:			
Patient Name:		Date:		

Mr BS DHEANSA FRCS MSc FRCS(Plast)
Consultant Plastic, Reconstructive & Cosmetic Surgeon

Telephone: 01342 330 383 Facsimile: 01342 330 301

Email: enquiry@my-plastic-surgeon.co.uk Website: www.my-plastic-surgeon.co.uk



ADDITIONAL NOTES OR COMMENTS:

ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
ı	
1	

Telephone: 01342 330 383 Facsimile: 01342 330 301 Email: enquiry@my-plastic-surgeon.co.uk Website: www.my-plastic-surgeon.co.uk Website: www.my-plastic-surgeon.co.uk